

V CONVEGNO AISSA #UNDER40

LE SCIENZE AGRARIE NELL'ANTROPOCENE:
DALLA PRODUTTIVITÀ ALLA TUTELA
DEL PATRIMONIO MATERIALE E CULTURALE

BOOK OF ABSTRACTS

26-27 GIUGNO 2024
UNIVERSITÀ DI FIRENZE,
CAMPUS DI NOVOLI, EDIFICIO D6

110 anni di
Agraria

Il futuro è
nell'agrAria



Con il patrocinio di:



Organizzato da:



UNIVERSITÀ
DEGLI STUDI
DI FIRENZE
DAGRI
DIPARTIMENTO DI SCIENZE
E TECNICHE AGRARIE,
ALIMENTARI, AMBIENTALI E FORESTALI

#AISSA
under40



Technological affinity index for interaction between lactic acid bacteria and *Saccharomyces cerevisiae* strains to modulate the fruity and floreal aroma of Catarratto wine

Vincenzo Naselli^{1*}, Antonino Pirrone¹, Enrico Viola¹, Venera Seminerio¹, Valentina Craparo¹, Micaela Carusi¹, Davide Alongi¹, Antonella Porrello², Antonella Maggio², Antonio Alfonzo¹, Luca Settanni¹, Giancarlo Moschetti¹, Nicola Francesca¹

¹Department of Agricultural, Food and Forest Sciences (SAAF), University of Palermo, Viale Delle Scienze Bldg. 5, Ent. C, 90128, Palermo, Italy

²Department of Biological, Chemical and Pharmaceutical Sciences and Technologies (STEBICEF), University of Palermo, Viale Delle Scienze, Building 17 Parco d'Orleans II, 90128, Palermo, Italy

*E-mail: vincenzo.naselli@unipa.it

Abstract

The sensory quality of wine is significantly influenced by microbial interactions. Specifically, the time (t), rate (m), and volatile organic compounds (VOC) associated with malolactic fermentation are linked to the interaction between yeast and lactic acid bacteria. The study investigated the interactions between *Lactiplantibacillus plantarum* or *Oenococcus oeni* and *Saccharomyces cerevisiae* by using the Technological Affinity Index (TAIndex) to assess these interactions. Interestingly, the parameter "m" exhibited a negative correlation with both TAIndex values and the duration of malolactic fermentation. The microbial associations observed in different trials were as follows: in the CO5 trial, where *L. plantarum* MLPK45H was coupled with *S. cerevisiae* NF213, the TAIndex value was 0.351, and the 'm' value was -0.65 (g/L×d⁻¹), while in the CO10 trial, the combination of *L. plantarum* MLPK45H and *S. cerevisiae* QA23 resulted in a TAIndex value of 0.348 and an 'm' value of -0.66 (g/L×d⁻¹). Under the provided conditions, malolactic fermentation occurred within two days. Furthermore, the CO1 trial involved consociations between the *O. oeni* MLB6 and *S. cerevisiae* NF213. In this case, the TAIndex value was 0.009, the "m" value of -0.0781 (g/L×d⁻¹), and the malolactic fermentation process extended over 44 days. This prolonged fermentation produced increased levels of ethyl octanoate and ethyl decanoate. The fruity and floral sensory perceptions were increased in CO1, probably due to lower concentrations of 3-methyl-1-butanol, hexanoic acid, octanoic acid and decanoic acid compared to the CONT A1 study, as evaluated by the panel.

In conclusion, the LAB-*S. cerevisiae* microbial consociation offers a valuable microbial approach to produce wines that are highly appreciated by panelists. The time and rate of malolactic fermentation significantly influence the synthesis of VOC with a pronounced olfactory impact. Consequently, knowledge of TAIndex could prove pivotal in enhancing oenological planning to produce wines with exceptionally intense fruity and floral characteristics.

Keywords

Lactic acid bacteria, Saccharomyces cerevisiae, malolactic fermentation, wine, technological affinity index